

Critically-damped oscillators and General Tau Theory exhibit similar error across speakers with different vocal tract dimensions

Christopher Geissler, Boston College
 Jyothiraditya Nellakra, University of Southern California
 geissle@bc.edu, nellakra@usc.edu

USC



Models


How do we describe the shape of an articulatory gesture?

- Theories of control & planning
 → Equations with known parameters
- “CDO”: Critically-Damped Oscillators
 - Mass-spring equation: $\ddot{x} + b\dot{x} + kx = 0$
 - $x(t) = D(e^{-\omega_0 t} + \omega_0 t e^{-\omega_0 t})$

“Tau”: General Tau Theory

- Gap-closure equation

$$x(t) = D \left(1 - \frac{t^2}{T^2}\right)^{\frac{1}{\kappa_{x,G}}}$$



Methods

- X-Ray Microbeam Dataset
 - Closure of C in [əCa]: [p b m f v t d n s z j ʒ]
 - Est. vocal tract length (VTL) from [ə] F3 & F4
- Unpack model equations to determine params
 - 2 additional parameters: x-offset and t-offset
 - CDO1 and Tau1: set parameters using GONS, PVEL, NONS. (5% vel. points + PVEL)
 - CDO2 and Tau2: set parameters using PVEL and first 20% velocity point.
 - CDO3 and Tau3: regress parameters to minimise error. (Theoretical best-fit)
- Error is mean Euclidean distance of modelled trajectory from experimental trajectory.
- Modelling and analysis was done using <High level summary of techniques and tools, details in next section>


References

Elie, Benjamin, David N. Lee & Alice Turk. 2023. Modeling trajectories of human speech articulators using general Tau theory. *Speech Communication* 151. 24–38. <https://doi.org/10.1016/j.specom.2023.04.004>.
 Lee, David N. 1998. Guiding Movement by Coupling Taus. *Ecological Psychology* 10(3/4). 221. <https://doi.org/10.1080/10407413.1998.9652683>.
 Saltzman, Elliot L. & Kevin G. Munhall. 1989. A Dynamical Approach to Gestural Patterning in Speech Production. *Ecological Psychology* 1(4). 333–382. https://doi.org/10.1207/s15326969eco0104_2.
 Westbury, John R., Greg Turner & J. Dembowski. 1994. *X-ray microbeam speech production database user's handbook, version 1.0*. University of Wisconsin Waisman Center.

Analysis & Results


Sample fits

Sample simulations




CDO results

CDO distance scales with magnitude error ~ magnitude + manner + place




Vocal tract results

No effect of VTL/sex/height, if use magnitude → oral cavity height matters more than VTL



Tau results

Tau fits better than CDO (note y-axis scaling) error ~ magnitude + manner + duration



Unpacking Models

- Regressing parameters for best-fit cannot account for implicit statements of models:
 - Acceleration at PVEL should be 0.
 → $\ddot{x}(t_{PVEL}) = 0$
 - Velocity at GONS and NONS should be ≈ 0.
 → $\dot{x}(t_{GONS}) \approx \dot{x}(t_{NONS}) \approx 0$
 - The velocity at the 20% points should be 1/3 of PVEL.
 → $\dot{x}(t_{20\%point}) = 0.2\dot{x}(t_{PVEL})$
- (a & b) → CDO1, Tau1. (a & c) → CDO2, Tau2
- x-offset and t-offset are also parameters.
 - Models treat start as $t = 0$ and end at $x = 0$.
 - Experimental data has arbitrary start times.
 - Goal might not be constriction degree $CD = 0$.
 - Conversion function requires x and t offset.
- Implications for a model of speech production:
 - Gesture execution requires two or three points (x, vel, acc, t). x is CD on a linear scale, $t = 0$ is arbitrary, depending on the gestural score.
 - Parameter setting for model $x(t)$ is automatic.
 - Model $x(t)$ function governs motor output.

Conclusions

- These models fit similarly across individuals
- New method of calculating model parameters
 - Below: Natural Frequency, Kappa
- Further work is needed to extend this analysis to models of gradient gestural activation over time

